

Supporting Resources

Use the links below to jump to a specific resource.

Six Types of Concussion

Information on the six types of clinical trajectories of concussion

A Fact Sheet for School Nurses

CDC's fact sheet specific to school nurses

Tennessee's Return to Learn / Return to Play: Concussion Management Guidelines

Good information about steps to return a child to the classroom and steps for a gradual return to play

CDC Online Training for Healthcare Providers

Earn free CME, CNE, and CEU credits

Research Summary and References

Summary of research provided in training and references

Changes to Watch for Over Time



CONCUSSION CLINICAL TRAJECTORIES

A Model for Understanding
Assessment, Treatment
and Rehabilitation

COGNITIVE/FATIGUE



Cognitive difficulties include decreased concentration, increased distractibility, difficulty learning/retaining new information or decreased multitasking abilities. Sometimes accompanied by increased fatigue as the day progresses.

VESTIBULAR



Impairments of the vestibular system - the balance center of the brain - affect one's ability to interpret motion, coordinate head and eye movements, or stabilize vision upon head movement.

OCULAR



Ocular dysfunction occurs when the movement of the eyes in tandem, or binocular eye movement, is affected. This may result in difficulties bringing the eyes together, or moving one's eyes to track motion.

POST-TRAUMATIC MIGRAINE




Post-traumatic migraine symptoms include headaches, nausea, and/or sensitivity to light or noise.

CERVICAL



Sometimes, the concussive blow affects the extra-cranial region including the neck and/or spinal cord. An injury of this type may lead to ongoing headaches.

ANXIETY/MOOD



This occurs when someone has a hard time turning his or her thoughts off, being particularly ruminative, of suffering from excessive worry or concern.

SIX TYPES OF CONCUSSION

- 1 Cognitive/Fatigue
- 2 Vestibular
- 3 Ocular
- 4 Post-traumatic Migraine
- 5 Cervical
- 6 Anxiety/Mood

FACTS

- ✦ Symptoms will be broad and generalized during the first week following concussion and will generally include symptoms like headache and fatigue.
- ✦ After the first week, if symptoms persist, they will tend to fall into one of the 6 clinical trajectories.
- ✦ There could be more than one trajectory type present.
- ✦ Specific trajectory and outcome depends on several factors:
 - Direction of force (linear vs. rotational)
 - Location of impact
 - Amount of force involved
 - Pre-injury risk factors

ACTIVE TREATMENT

Research is showing that active, specialized treatment – focused on specific symptoms – helps the brain recover from injury.

- ✦ Neuropsychology
- ✦ Vestibular Physical Therapy
- ✦ Exertional Physical Therapy
- ✦ Physical Medicine and Rehabilitation
- ✦ Neuro-optometry/ Neuro-ophthalmology
- ✦ Orthopedist
- ✦ Neurosurgery
- ✦ Neuroradiology
- ✦ Chiropractic
- ✦ Cognitive Therapy/ Speech Language Pathology

RISK FACTORS (which may delay recovery)

- ✦ History of prior concussions
- ✦ Motion sickness
- ✦ Visual problems
- ✦ Learning or attention issues
- ✦ Migraine history
- ✦ Gender (*female*)
- ✦ Age (*younger children tend to take longer to recover*)

Source: Kontos, A.P. Collins, M.W., (2018). *Concussion: A Clinical Profile Approach to Assessment and Treatment*.



A Fact Sheet for School Nurses

**HEADS UP
SCHOOLS**

THE FACTS:

- * All concussions are serious.
- * Most concussions occur without loss of consciousness.
- * Recognition and proper response to concussions when they first occur can help aid recovery and prevent further injury, or even death.

What is a concussion?

A concussion is a type of brain injury that changes the way the brain normally works. A concussion is caused by a bump, blow, or jolt to the head. Concussions can also occur from a fall or blow to the body that causes the head and brain to move rapidly back and forth. Even what seems to be a mild bump to the head can be serious.

How can I recognize a concussion?

To help you recognize a concussion, ask the injured student or witnesses of the incident about:

1. Any kind of forceful blow to the head or to the body that resulted in rapid movement of the head.

-and-

2. Any change in the student's behavior, thinking, or physical functioning. (See the signs and symptoms of concussion.)

To download this fact sheet in Spanish, please visit: www.cdc.gov/Concussion.
Para obtener una copia electrónica de esta hoja de información en español, por favor visite: www.cdc.gov/Concussion.

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION



How can concussions happen in schools?

Children and adolescents are among those at greatest risk for concussion. Concussions can result from a fall, or any time a student's head comes into contact with a hard object, such as the floor, a desk, or another student's head or body. The potential for a concussion is greatest during activities where collisions can occur, such as during physical education (PE) class, playground time, or school-based sports activities.

Students may also get a concussion when doing activities outside of school, but then come to school when symptoms of the concussion are presenting. For example, adolescent drivers are at increased risk for concussion from motor vehicle crashes.

Concussions can have a more serious effect on a young, developing brain and need to be addressed correctly. Proper recognition and response to concussion symptoms in the school environment can prevent further injury and can help with recovery.



What are the signs and symptoms of concussion?

Students who experience **one or more** of the signs and symptoms listed below after a bump, blow, or jolt to the head or body should be referred to a health care professional experienced in evaluating for concussion.

There is no one single indicator for concussion. Rather, recognizing a concussion requires a symptom assessment. The signs and symptoms of concussion can take time to appear and can become more noticeable during concentration and learning activities in the classroom. For this reason, it is important to watch for changes in how the student is acting or feeling, if symptoms become worse, or if the student just “doesn't feel right.”

SIGNS OBSERVED BY SCHOOL NURSES

- Appears dazed or stunned
- Is confused about events
- Answers questions slowly
- Repeats questions
- Can't recall events *prior* to the hit, bump, or fall
- Can't recall events *after* the hit, bump, or fall
- Loses consciousness (even briefly)
- Shows behavior or personality changes

SYMPTOMS REPORTED BY THE STUDENT

Thinking/Remembering:

- Difficulty thinking clearly
- Difficulty concentrating or remembering
- Feeling more slowed down
- Feeling sluggish, hazy, foggy, or groggy

Emotional:

- Irritable
- Sad
- More emotional than usual
- Nervous

Physical:

- Headache or “pressure” in head
- Nausea or vomiting
- Balance problems or dizziness
- Fatigue or feeling tired
- Blurry or double vision
- Sensitivity to light or noise
- Numbness or tingling
- Does not “feel right”

Sleep*:

- Drowsy
- Sleeps *less* than usual
- Sleeps *more* than usual
- Has trouble falling asleep

**Only ask about sleep symptoms if the injury occurred on a prior day.*



Remember, you can't see a concussion and some students may not experience or report symptoms until hours or days after the injury. Most young people with a concussion will recover quickly and fully. But for some, concussion signs and symptoms can last for days, weeks, or longer.



What are concussion danger signs?

In rare cases, a dangerous blood clot may form on the brain in a person with a concussion and crowd the brain against the skull. The student should be taken to an emergency department right away if s/he exhibits any of the following danger signs after a bump, blow, or jolt to the head or body:

- One pupil larger than the other
- Is drowsy or cannot be awakened
- A headache that gets worse and does not go away
- Weakness, numbness, or decreased coordination

- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Cannot recognize people or places
- Becomes increasingly confused, restless, or agitated
- Has unusual behavior
- Loses consciousness (even a brief loss of consciousness should be taken seriously)

For more information and tool kits for youth sports coaches and high school coaches, visit www.cdc.gov/Concussion.

What can school nurses and school professionals do?

Below are steps for you to take when a student comes to your office after a bump, blow, or jolt to the head or body.

1. **Observe student for signs and symptoms of concussion for a minimum of 30 minutes.**
2. **Complete the *Concussion Signs and Symptoms Checklist* and monitor students consistently during the observation period.** The form includes an easy-to-use checklist of signs and symptoms that you can look for when the student first arrives at your office, fifteen minutes later, and at the end of 30 minutes, to determine whether any concussion symptoms appear or change.
3. **Notify the student's parent(s) or guardian(s) that their child had an injury to the head.**
 - > **If signs or symptoms are present:** refer the student right away to a health care professional with experience in evaluating for concussion. Send a copy of the *Concussion Signs and Symptoms Checklist* with the student for the health care professional to review. Students should follow their health care professional's guidance about when they can return to school and to physical activity.

- > **If signs or symptoms are not present:** the student may return to class, but should not return to sports or recreation activities on the day of the injury. Send a copy of the *Concussion Signs and Symptoms Checklist* with the student for their parent(s) or guardian(s) to review and ask them to continue to observe the student at home for any changes. Explain that signs and symptoms of concussion can take time to appear. Note that if signs or symptoms appear, the student should be seen right away by a health care professional with experience in evaluating for concussion.



Children and teens with a concussion should NEVER return to sports or recreation activities on the same day the injury occurred. They should delay returning to their activities until a health care professional experienced in evaluating for concussion says they are symptom-free and it's OK to return to play. This means, until permitted, not returning to:

- Physical Education (PE) class,
- Sports practices or games, or
- Physical activity at recess.



What do I need to know about students returning to school after a concussion?

Supporting a student recovering from a concussion requires a collaborative approach among school professionals, health care professionals, parents, and students. All school staff, such as teachers, school nurses, counselors, administrators, speech-language pathologists, coaches, and others should be informed about a returning student's injury and symptoms, as they can assist with the transition process and making accommodations for a student. If symptoms persist, a 504 meeting may be called. Section 504 Plans are implemented when students have a disability (temporary or permanent) that affects their performance in any manner. Services and accommodations for students may include speech-language therapy, environmental



School Policies:

Students Returning to School after a Concussion

Check with your school administrators to see if your district or school has a policy in place to help students recovering from a concussion succeed when they return to school. If not, consider working with your school administration to develop such a policy. Policy statements can include the district's or school's commitment to safety, a brief description of concussion, a plan to help students ease back into school life (learning, social activity, etc.), and information on when students can safely return to physical activity following a concussion.

adaptations, curriculum modifications, and behavioral strategies.

Encourage teachers and coaches to monitor students who return to school after a concussion. Students may need to limit activities while they are recovering from a concussion. Exercising or activities that involve a lot of concentration, such as studying, working on the computer, or playing video games, may cause concussion symptoms (such as headache or tiredness) to reappear or get worse. After a concussion, physical and cognitive activities—such as concentration and learning—should be carefully monitored and managed by health and school professionals.

If a student already had a medical condition at the time of the concussion (such as chronic headaches), it may take longer to

recover from the concussion. Anxiety and depression may also make it harder to adjust to the symptoms of a concussion.

School professionals should watch for students who show increased problems paying attention, problems remembering or learning new information, inappropriate or impulsive behavior during class, greater irritability, less ability to cope with stress, or difficulty organizing tasks. Students who return to school after a concussion may need to:

- Take rest breaks as needed,
- Spend fewer hours at school,
- Be given more time to take tests or complete assignments,
- Receive help with schoolwork, and/or
- Reduce time spent on the computer, reading, or writing.

It is normal for a student to feel frustrated, sad, and even angry because s/he cannot return to recreation or sports right away, or cannot keep up with schoolwork. A student may also feel isolated from peers and social networks. Talk with the student about these issues and offer support and encouragement. As the student's symptoms decrease, the extra help or support can be gradually removed.

What can I do to prevent and prepare for a concussion?

Here are some steps you can take to prevent concussions in school and ensure the best outcome for your students:

Prepare a concussion action plan. To ensure that concussions are identified early and managed correctly, have an action plan in place before the start of the school year. This plan can be included in your school or district's concussion policy. You can use the online action plan for sports and recreation activities at: www.cdc.gov/concussion/response/html. Be sure that other appropriate school and athletic staff know about the plan and have been trained to use it.

Educate parents, teachers, coaches, and students about concussion. Parents, teachers, and coaches know their students well and may be the first to notice when a student is not acting normally. Encourage teachers, coaches, and students to:

- Learn about the potential long-term effects of concussion and the dangers of returning to activity too soon.
- Look out for the signs and symptoms of concussion and send students to see you if they observe any or even suspect that a concussion has occurred.
- View videos about concussion online at: www.cdc.gov/Concussion.

Prevent long-term problems. A repeat concussion that occurs before the brain recovers from the previous concussion—usually within a short period of time (hours, days, or weeks)—can slow recovery or increase the likelihood of having long-term problems. In rare cases, repeat concussions

can result in edema (brain swelling), permanent brain damage, and even death. Keep students with a known or suspected concussion out of physical activity, sports, or playground activity on the day of the injury and until a health care professional with experience in evaluating for concussion says they are symptom-free and it is OK for the student to return to play.

Create safe school environments.

The best way to protect students from concussions is to prevent concussions from happening. Make sure your school has policies and procedures to ensure that the environment is a safe, healthy place for students. Talk to all school staff and administrators and encourage them to keep the physical space safe, keep stairs and hallways clear of clutter, secure rugs to the floor, and check the surfaces of all areas where students are physically active, such as playing fields and playgrounds. Playground surfaces should be made of shock-absorbing material, such as hardwood mulch or sand, and maintained to an appropriate depth. Proper supervision of students is also important.



For more detailed information about concussion diagnosis and management, please download *Heads Up: Facts for Physicians about Mild Traumatic Brain Injury* from CDC at: www.cdc.gov/Concussion.

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION



Monitor the health of your student athletes.

Make sure to ask whether an athlete has ever had a concussion and insist that your athletes are medically

evaluated and are in good condition to participate in sports. Keep track of athletes who sustain concussions during the school year. This will help in monitoring injured athletes who participate in multiple sports throughout the school year.

Some schools conduct preseason baseline testing (also known as neurocognitive tests) to assess brain function—learning and memory skills, ability to pay attention or concentrate, and how quickly someone can think and solve problems. If an athlete has a concussion, these tests can be used again during the season to help identify the effects of the injury. Before the first practice, determine whether your school would consider baseline testing.

Again, remember your
concussion ABCs:

- A—Assess the situation
- B—Be alert for signs and symptoms
- C—Contact a health care professional

HEADS UP
SCHOOLS

* For more information on concussion and to order additional materials for school professionals **FREE-OF-CHARGE**, visit: www.cdc.gov/Concussion.



Return to Learn/Return to Play: Concussion Management Guidelines

Tennessee Department of Health | August 2020



Acknowledgments

June, 2017: Rev. Nov, 2019: Aug, 2020

This document is a compilation of concussion management material produced by the states of Colorado and Nebraska and has been adapted with permission for use by the Tennessee Department of Health.

This document has been created/edited/reviewed by the following people:

Tim Lee, MHA, MS, ATC, Vanderbilt Sports Concussion Center

Alex Diamond, DO, MPH, Vanderbilt Sports Medicine/Vanderbilt Sports Concussion Center/Program for Injury Prevention in Youth Sports

Gary Solomon, PhD, Clinical Neuropsychologist, Vanderbilt Sports Concussion Center

Paula Denslow, CBIS, Director, Brain Links, Tennessee Disability Coalition

Wendy Ellmo, MS, CCC-SLP, BCNCDS, Brain Links, Tennessee Disability Coalition

Lori Paisley, Executive Director, Coordinated School Health, TN Department of Education

Dereese Methvin, EdS, RN, BSN, Coordinated School Health, TN Department of Education

Andrew Gregory, MD, Vanderbilt Sports Concussion Center

Carolina Clark, MD, MPH, Child Health Medical Consultant, TN Department of Health

Morgan F. McDonald, MD, FAAP, FACP, Assistant Commissioner, Director, Division of Family Health and Wellness, TN Department of Health

Michelle D. Fiscus, MD, FAAP, Deputy Medical Director, Chronic Disease and Injury Prevention, Health Promotion, TN Department of Health

Rachel Heitmann, MS, Section Chief, Injury Prevention and Detection TN Department of Health

Jean Doster, MS, Past Director, Traumatic Brain Injury Program, TN Department of Health

Ashley Bridgman, MS, Director, Traumatic Brain Injury Program, TN Department of Health

This document can be viewed online at <https://www.tn.gov/health/health-program-areas/fhw/vipp/tbi/resources.html>



Tennessee Department of Health Authorization Number 343602. This public document was promulgated at a cost of \$0.00 per copy. 0 copies 08-2020

Table of Contents

Section 1: Education

What is a Concussion/TBI?	4
Why are Concussions/TBI a Big Deal?	5
Signs and Symptoms of Concussion	6
Prevention	7

Section 2: Return to Learning

Concussion Management Team	8
The Concussion Management Process (Example)	10
Returning to School	11
Classroom Strategies for Concussion Recovery	12
Symptoms Checklist	13
When and How to Write a 504 Plan	14

Section 3: Return to Play

Law Return to Play	15
Tennessee State Sports Concussion	15
Law Return to Play Decisions	16
Graduated Return to Play Plan	16

Section 4: References 17

Section 5: Additional Resources 17

Section 6: Appendices

A. Cognitive Activity Monitoring Log	18
B. Authorization of Release of Medical Information	19
C. School Accommodations	20
D. Return to Play	21 - 22

What is a Concussion?

Concussion/TBI

A concussion is a type of traumatic brain injury, or TBI, is caused by a bump, blow or jolt to the head or by a hit to the body that causes the head and brain to move rapidly back and forth. This sudden movement can cause the brain to bounce around or twist in the skull, creating changes in the brain, and sometimes stretching and damaging the brain cells (CDC, 2015).

Aside from the elderly, children and adolescents are among those at greatest risk for concussion. The potential for a concussion in young people is greatest during activities where collisions can occur, such as during physical education class, playground time or sports activities. However, concussions can happen any time a student's head comes into contact forcefully with a hard object, such as a floor, desk or another student's head or body. Proper recognition and response to concussion can prevent further injury and help with recovery (CDC, 2015).

Medical providers may describe a concussion as a "mild" brain injury because concussions are usually not life-threatening. Even so, the effects of a concussion can be serious (CDC, 2015).

Traumatic brain injury is a serious public health problem in the United States. Each year, traumatic brain injuries contribute to a substantial number of deaths and cases of permanent disability. In 2014, 2.5 million TBIs occurred either as an isolated injury or along with other injuries (CDC, 2015).



Why are Concussions a Big Deal?

A concussion can occur from an impact to the body or the head. The most common cause of a concussion is a whiplash type injury, involving a rapid acceleration of the head.

Most concussions (90 percent) occur without loss of consciousness. Concussions can occur in any sport or during regular daily activities.

A “ding,” “getting your bell rung” or what seems to be a mild bump, blow or jolt to the head can be serious and can change the way the brain normally works (CDC, 2013).

Because of changes in the neurophysiology of the brain, symptoms may continue to develop over the next few days following an injury.

After a concussion, among other effects, nerve cells and connections within the brain become stressed, resulting in the possible breaking of some connections between different brain areas and limiting the ability of the brain to process information efficiently and quickly (Molfese, 2013).

These changes can lead to a set of symptoms affecting the student’s cognitive, physical, emotional and sleep functions, which may result in reduced ability to do tasks at home, at school or at work. Concussions can have an impact on the student’s ability to learn in the classroom. Tracking symptoms tells a big part of the story during recovery.

During this time of recovery, returning to play before symptoms have resolved incurs the risk of further injury, and returning to full-time academics before symptoms have cleared can result in prolonged recovery time.

As the chemistry of the brain returns to normal, the symptoms begin to subside and for most people, they resolve within one to four weeks. During the recovery period, monitor students for full resolution of symptoms and refer for further evaluation or treatment if needed.

Ignoring the symptoms and trying to “tough it out” often makes symptoms worse.

Second Impact Syndrome may occur when a brain already injured takes another blow or hit before the brain recovers from the first, usually within a short period of time (hours, days or weeks). A repeat concussion can slow recovery or increase the likelihood of having long-term problems. In rare cases, repeat concussions can result in edema (brain swelling), permanent brain damage and even death (CDC, 2013).

(Adapted from Return to Learn, 2014)

Signs and Symptoms of Concussions

The signs and symptoms of concussion can show up right after an injury or may not appear or be noticed until hours or a few days after the injury. Be alert for any of the following signs or symptoms. Also, watch for changes in how the student is acting or feeling, if symptoms are getting worse or if the student just "doesn't feel right" (CDC, 2015).

Signs Reported by the Student:

Emotional:

- Irritability
- Sadness
- More emotional than usual
- Nervousness

Physical:

- Headache or "pressure" in head
- Nausea or vomiting
- Balance problems or dizziness
- Fatigue or feeling tired
- Blurry or double vision
- Numbness or tingling
- Does not "feel right"

Signs observed by staff:

- Appears dazed or stunned
- Is confused about events
- Answers questions slowly
- Repeats questions
- Can't recall events prior to the hit, bump or fall
- Can't recall events after the hit, bump or fall
- Loses consciousness (even briefly)
- Shows behavior or personality changes
- Forgets class schedule or assignments

Cognitive:

- Difficulty thinking clearly
- Difficulty remembering or concentrating
- Feeling slowed down
- Feeling sluggish, hazy or foggy

Sleep:

- Drowsy
- Sleeps less than usual
- Sleeps more than usual
- Has trouble falling asleep (Only ask sleep symptoms if injury occurred prior to date reported)

Danger Signs:

Be alert for symptoms that worsen over time. A student should be seen in the emergency department right away if s/he has:

- **One pupil that is larger than the other**
- **Drowsiness or cannot be awakened**
- **A headache that gets worse and does not go away**
- **Weakness, numbness or decreased coordination**
- **Repeated vomiting**
- **Slurred speech**
- **Seizures**
- **Difficulty recognizing people or places**
- **Increased confusion, restlessness or agitation**
- **Unusual behavior**
- **Loss of consciousness**

Prevention

A concussion is a traumatic brain injury that can be prevented in many cases. Being an active participant in sports and engaging in physical activity does place student-athletes at higher risk for injury; however, there are preventive measures that schools can take. This section is intended to remind school districts about the importance of prevention. Schools should:

- Conduct periodic safety reviews of common play/sporting areas
- Provide appropriate and adequate staffing for sporting events and recess
- Provide appropriate access to protective gear (helmets, mouth guards)
- Provide appropriate fitting of protective gear
- Design guidelines and enforcement of appropriate and fair rules and techniques (CDE, 2014)

Design, Implement and Review a school-wide “concussion action plan” for all school staff and faculty. Know what to do BEFORE a student/athlete has an injury.

Implement Safe Stars Initiative

The Safe Stars initiative recognizes youth sports leagues throughout Tennessee for providing the highest level of safety for their youth athletes. Safe Stars consists of three levels: gold, silver and bronze, and involves implementation of policies around topics such as concussion education, weather safety and injury prevention.

Safe Stars’ goal is to provide resources and opportunities for every youth sports league to enhance their safety standards. The criteria for achieving recognition as a Safe Stars league has been developed by a committee of health professionals dedicated to reducing sports-related injuries among youth.

To learn more please visit:


www.tn.gov/health/health-program-areas/fhw/vipp/safe-stars-initiative.html.



Concussion Management Team

Once a concussion has been diagnosed by a health care professional, managing the concussion is best accomplished by creating a support system for the student. Communication and collaboration among parents, school personnel, coaches, athletic trainers and health care providers is essential for the recovery process. This support system oversees the return to academics and return to play process. A medical release signed by the parents allows for two-way communication between the school personnel and the health care provider (McAvoy, 2012, Return to Learn, 2014).

A collaborative approach with the student as the focus!



Each school district creates a concussion management policy that incorporates:

- Knowledge about concussions as a mild traumatic brain injury
- Training for all coaches, athletes, parents and school staff members about concussion management
- A Concussion Management Team with a designated Concussion Management Team Point Person
 - o The Concussion Management Point Person may be the school nurse, the 504 designee, a guidance counselor or an administrator. Choose the individual that works best for your school's situation.

The Concussion Management Team

Members may include:

Physicians

Speech Language

Neuropsychologists

Pathologist Nurse

Physician Assistant

Practitioner

Parents

School Nurse

School Administrator or

School Psychologist

Designee

School Counselor

Athletic Director

Occupational Therapist

Athletic Trainer

Physical Therapist

Coach

Student-Athlete

Teacher


(Return to Play, 2014)



TEAMWORK

The Concussion Management Process

This is an example of the concussion management process that includes best practice components for all students.



(Adapted from Colorado, 2014)

Returning to School

The student may return to school when symptoms are tolerable and manageable, **as long as the school is making appropriate accommodations for the student**. The school must understand concussions and the necessary academic accommodations in order to facilitate returning students to the learning environment.


Key points:

- If symptoms prevent the student from concentrating on mental activities for ten minutes or less, complete cognitive rest is required. The student should be kept home from school with limited external stimulation (texting, watching TV, playing video games, etc.) or driving. In some, but not all, cases these stimulating activities may worsen the symptoms of concussion.
- If symptoms allow the student to concentrate on mental activities for up to 20 minutes or less, parents should consider keeping the student home from school, but may allow increased time periods of external stimulation as long as symptoms do not get worse.
- **See Cognitive Activity Monitoring Log in Appendix A**

When the student can tolerate 30 minutes of light mental activity, parents can consider returning him or her to the classroom. Best practices suggest: (a) parents communicate with the school and sign **a medical release of information (See Appendix B)** for the school to communicate with the health care provider, and (b) implement the appropriate academic accommodations provided by the treating health care provider and concussion management team.

Academic Accommodations: See School Accommodations Template in Appendix C

The balance between the student's medical and academic needs should be closely coordinated between school personnel and the health care provider. Each concussed student can have different symptoms, a different level of severity and a different recovery. Academic accommodations should be tailored to the specific needs of the individual student (McAvoy, 2014). Certain symptoms lend themselves to certain interventions. Especially in the acute phase of the concussion (one-four weeks), interventions should be applied generously in the classroom setting. Symptoms may be worse in some classes than in others. Teachers are encouraged to apply any intervention that is needed for the student based on the symptoms (McAvoy, 2015).



Classroom Strategies for Concussion Recovery

Symptom	School Setting Adjustment
Headache	<ul style="list-style-type: none"> • Frequent breaks • Reduce exposure to specific aggravators: brightlights/computer work/noisy environment • Rest periods if needed in nurse's office or quiet environment
Dizziness	<ul style="list-style-type: none"> • Allow student to put head down on desk • Give student early dismissal from class to avoid crowded hallways
Visual Problems: Light Sensitivity, Double Vision, Blurry Vision	<ul style="list-style-type: none"> • Reduce exposure to computers, light boards, videos • Reduce brightness on screens • Allow student to wear hat/sunglasses • Consider use of audio books • Turn off fluorescent lights • Seat student closer to the center of the classroom (blurryvision) • Have school nurse cover one eye with a patch for students with double vision
Noise Sensitivity	<ul style="list-style-type: none"> • Allow student to have lunch in a quiet area with one classmate • Limit/avoid band, choir, shop classes • Consider use of ear plugs • Allow early dismissal from classto avoid noisy hallways • Avoid noisy gyms/sporting events
Difficulty Concentrating or Remembering	<ul style="list-style-type: none"> • Avoid testing or completingmajor projects during recovery • Allow extra time to complete non-standardized tests • Postpone standardized testing • Consider one test per dayduring exams • Consider use of notes, a note taker or reader for oral testing
Sleep Disturbance	<ul style="list-style-type: none"> • Allow for late start or short dayto catch up on sleep • Allow rest breaks in a quiet area

Adapted from: Halstead, M.E., McAvoy, K., Devore, C.D., Carl, R., Lee, M., Logan, K. (2013). Return to learning following a concussion. *American Academy of Pediatrics*. 132: 5, 948-957.doi:10.1542/peds.2013-2867

Symptoms Checklist

In most cases, symptoms may be the primary way to know when and how a concussion is getting better. Since the report of symptoms can be quite subjective, it is helpful to use a rating scale. The rating scale can act as a common language for everyone involved in managing the concussion. Most concussion management programs utilize a symptom scale with a 0 to 6 rating scale (0 = not present; 6 = most severe).

Name: _____ Date: _____

Date of Injury: _____

Symptom	None	Mild		Moderate		Severe	
Headache	0	1	2	3	4	5	6
Nausea	0	1	2	3	4	5	6
Vomiting	0	1	2	3	4	5	6
Balance problems	0	1	2	3	4	5	6
Dizziness	0	1	2	3	4	5	6
Fatigue	0	1	2	3	4	5	6
Trouble falling asleep	0	1	2	3	4	5	6
Sleeping more than usual	0	1	2	3	4	5	6
Sleeping less than usual	0	1	2	3	4	5	6
Drowsiness	0	1	2	3	4	5	6
Sensitive to light	0	1	2	3	4	5	6
Sensitive to noise	0	1	2	3	4	5	6
Irritability	0	1	2	3	4	5	6
Sadness	0	1	2	3	4	5	6
Nervous/Anxious	0	1	2	3	4	5	6
Feeling more emotional	0	1	2	3	4	5	6
Numbness or tingling	0	1	2	3	4	5	6
Feeling like in a fog	0	1	2	3	4	5	6
Difficulty remembering	0	1	2	3	4	5	6
Difficulty concentrating	0	1	2	3	4	5	6
Visual problems	0	1	2	3	4	5	6
Total Symptoms Score							

The Graded Symptoms Checklist is recommended by the National Athletic Trainers Association (Casa et al., 2012). The 0 to 6 symptoms scale is commonly used by various tests: ImPACT and SCAT3.

(Adapted from Colorado, 2014)

When and How to Write a 504 Plan

Typically, 90 percent of kids with concussions will recover within four weeks of their injuries. If a student has not resolved from a concussion within the typical three to four week time frame, it may be prudent to begin to look at a more “targeted” approach. (McAvoy and Eagan, 2015). If a 504 Plan is indicated, the 504 designee (CMT Point Person) at the school should set up a meeting with all the necessary members of the concussion management team (teachers, parents, counselors, administrators, school nurse, etc.). When writing a 504 Plan, one must identify what the most problematic symptoms are which will let you know which interventions to use in your plan. There are certain conditions or “modifiers” of concussion that we know may prolong the recovery process. Those modifiers are:

- A history of migraine headache or family history of migraines
- A pre-existing headache disorder
- ADHD
- A history of previous concussions
- Learning disability
- A history of anxiety and depression
- Sleep disorder

Be specific in the writing you 504 Plan. Do not write a plan “for concussion”; use the phrasing, “Section 504 Plan for X (specified symptom) secondary to concussion.

Examples:

<p>Section 504 Plan for Headaches secondary to a concussion</p>	<ul style="list-style-type: none"> • Head down on the desk in classroom • Pass to leave room to visit nurse • Able to take medications in school clinic
<p>Section 504 Plan for Slowed Processing Speed secondary to a concussion</p>	<p>Appropriate Interventions:</p> <ul style="list-style-type: none"> • Extended time on in-class assignments • Extended time on tests
<p>Section 504 Plan for Convergence Insufficiency secondary to a concussion</p> <p>(MacAvoy & Eagan Brown, 2015)</p>	<p>Appropriate Interventions:</p> <ul style="list-style-type: none"> • Teacher or peer notes printed out • In-class and homework on paper instead of computer screens whenever possible • Books on tape

There should also be an overall medical and education plan addressing the following questions:

- How long do we expect the symptoms to linger?
- Is the student still being treated for his/her concussion/symptoms?
- Do we expect the student to fully recover?
- What are the medical interventions being used?
- What side effect should we expect?

Remember:

- Only a small percentage of students with a concussion will need a 504 Plan.
- A Release of Medical Information Form will be needed for the school to communicate with the medical provider (Appendix B).
- When the Concussion Management Team works together to identify the underlying cause(s) for the prolonged recovery, addresses those areas, supports the student with academic accommodations, monitors the progress and adjusts the plan as needed, full recovery is possible (McAvoy and Eagan- Brown, 2015).

Return to Play

Tennessee Sports Concussion Law

In April 2013, Tennessee became the 44th state to pass a sport concussion law designed to reduce youth sports concussions and increase awareness of traumatic brain injury.

The legislation, [Public Chapter 148](#), has three key components:

- To inform and educate coaches, youth athletes and their parents and require them to sign a concussion information form before competing.
- To require removal of a youth athlete who appears to have suffered a concussion from play or practice at the time of the suspected concussion.
- To require a youth athlete to be cleared by a licensed health care professional before returning to play or practice.

Both public and private school sports and recreational leagues for children under age 18 that require a fee are affected by the law. The law covers all sports. This website contains all the resources coaches, youth athletes and parents need to fulfill the intent of the law.

See more at:

<https://www.tn.gov/health/health-program-areas/fhw/vipp/tbi/tn-sports-concussion.html>

(TN Sports Concussion Law, 2013)

Within the school setting, any student who shows signs or symptoms of a concussion should be removed from physical activity (recess, physical education, dance class, etc.), and needs to be cleared medically before returning to physical activity. Medical providers approved to clear children for return to play from concussion are as follows:

- Medical Doctor (MD)
- Osteopathic Physician (DO)
- Clinical Neuropsychologist (PhD) with concussion training
- Physician Assistant (PA) with concussion training who is a member of a health care team supervised by a Tennessee licensed medical doctor or osteopathic physician.

See Return to Play Example, Appendix D

Return to Play Decisions

- ❑ According to the Concussion in Sport Group-4 Guidelines (2013), any child who is suspected of having a concussion should be removed from play and should not return to play that day.
- ❑ No return to sport should be considered until the child has returned to school successfully. A successful return to school would mean they no longer are in need of school accommodations.
- ❑ Children should not be returning to physical activity if they are still experiencing concussion symptoms, unless otherwise directed by their treating health care provider.
- ❑ Children should not be taking any medications to mask concussion symptoms in the graduated return to play process
- ❑ A graduated return to play process is recommended to be performed by the child with symptom monitoring at each step (McCrory, 2013).

Gradual Return to Play Plan

Return to play should occur in gradual steps beginning with light aerobic exercise only to increase your heart rate (e.g., stationary cycle); moving to increasing your heart rate with movement (e.g., running); then adding controlled contact if appropriate; and finally return to sports competition. Pay careful attention to your symptoms and your thinking and concentration skills at each stage or activity. After completion of each step without recurrence of symptoms, you can move to the next level of activity the next day under the direction of your health care provider. Move to the next level of activity only if you do not experience any symptoms at the present level. If your symptoms return, let your health care provider know, and await further instructions.

Day 1: Low levels of physical activity (i.e., symptoms do not come back during or after the activity). This includes walking, light jogging, light stationary biking and light weightlifting (low weight – moderate reps, no bench, no squats).

Day 2: Moderate levels of physical activity with body/head movement. This includes moderate jogging, brief running, moderate intensity on the stationary cycle, moderate intensity weightlifting (reduce time and or reduced weight from your typical routine).

Day 3: Heavy non-contact physical activity. This includes sprinting/running, high intensity stationary cycling, completing the regular lifting routine, non-contact sport specific drills (agility – with three planes of movement).

Day 4: Sports-specific practice.

Day 5: Full contact in a controlled drill or practice.

Day 6: Return to competition.

(TN Sports Concussion Law, 2013)

References:

1. Centers for Disease Control and Prevention-Concussion Fact Sheet for School Professionals (2013). www.cdc.gov/headsup/pdfs/custom/headsupconcussion_fact_sheet_for_schools.pdf
2. Centers for Disease Control and Prevention-Basic Information about Traumatic Brain Injury and Concussion. www.cdc.gov/traumaticbraininjury/basics.html
3. Colorado Department of Education Concussion Management Guideline (2014). www.cde.state.co.us/healthandwellness/concussionguidelines7-29-2014-0
4. Return to Learn: Bridging the Gap from Concussion to the Classroom (2014). www.education.ne.gov/sped/birsst/BRIDGING%20THE%20GAP%20Booklet%20plus%20Appendices.pdf
5. Halstead ME, McAvoy K, Devore CD, et al. Returning to learning following a concussion. *Pediatrics*. 2013;132(5):948-957. doi:10.1542/peds.2013-2867 <https://pubmed.ncbi.nlm.nih.gov/24163302/>
6. McAvoy, K & Eagan-Brown, B (2015). When to write a 504 Plan: Part 1. <http://nebula.wsimg.com/b7526f238f9c921d00a11d9a5769226f?AccessKeyId=E4B9300FA35CD0310DEE&disposition=0&alloworigin=1>
7. McAvoy, K & Eagan-Brown, B (2015). When to write a 504 Plan: Part 2. <http://nebula.wsimg.com/ce204367cd9b3c9ca304eceda7bbbb2c?AccessKeyId=E4B9300FA35CD0310DEE&disposition=0&alloworigin=1>
8. McAvoy, K & Eagan-Brown, B (2015). How to write a 504 Plan. <http://nebula.wsimg.com/b39dbbc04d4ee60fc623f9da5fedb363?AccessKeyId=E4B9300FA35CD0310DEE&disposition=0&alloworigin>
9. Tennessee Sports Concussion Law (2013). <https://www.tn.gov/health/health-program-areas/fhw/vipp/tbi/tn-sports-concussion.html>
10. McCorry, P., Meeuwisse, W. H., Aubry, M., Cantu, B., Dvořák, J., Echemendia, R. J., & Sills, A.(2013). Consensus statement on concussion in sport: the 4th International Conference on Concussion in Sport held in Zurich, November 2012. *British journal of sports medicine*, 47(5), 250-258.

Additional Resources:

1. Brain Links <http://tndisability.org/brain>
2. Center on Brain Injury Research & Training. <https://cbirt.org>
3. Colorado Kids with Brain Injury. <https://cokidswithbraininjury.com/>
4. Get Schooled on Concussions: Return to Learn. www.getschooledonconcussions.com/

Appendix A

Cognitive Activity Monitoring (CAM) Log

Name _____

Parent/ Teacher: _____

DATE TIME							
LOCATION (circle one)	Home School	Home School	Home School	Home School	Home School	Home School	Home School
COGNITIVE ACTIVITY:							
DURATION:							
SYMPTOM (PRE/POST)	Rate 0-10	Rate 0-10	Rate 0-10	Rate 0-10	Rate 0-10	Rate 0-10	Rate 0-10
HEADACHE	_/_	_/_	_/_	_/_	_/_	_/_	_/_
FATIGUE	_/_	_/_	_/_	_/_	_/_	_/_	_/_
CONCENTRATION PROBLEMS	_/_	_/_	_/_	_/_	_/_	_/_	_/_
IRRITABILITY	_/_	_/_	_/_	_/_	_/_	_/_	_/_
FOGGINESS	_/_	_/_	_/_	_/_	_/_	_/_	_/_
LIGHT/ NOISE SENSITIVITY	_/_	_/_	_/_	_/_	_/_	_/_	_/_
Other: _____	_/_	_/_	_/_	_/_	_/_	_/_	_/_
PRE-POST DIFFERENCE	_____	_____	_____	_____	_____	_____	_____

Appendix B

Authorization of Release of Medical Information for Concussion

School Name: _____
Patient Name: _____ Date of Birth: _____
Address: _____
City: _____ State: _____ Zip: _____
Social Security #: _____
I hereby authorize: _____
Name of Person/Organization Disclosing PHI

To release the following information to (School Receiving PHI) School: _____
Name: _____ Title: _____
Address: _____
Phone: _____ Fax: _____
Email: _____

Information to be shared:

- Medical records pertaining to concussion care
- Progress Notes
- Other: _____
- Academic Accommodations Forms
- Mental/Behavioral health records

The Information may be disclosed for the following purpose(s) only:

- Continued Treatment
- At the request of the patient/legal guardian

I understand that by voluntarily signing this authorization:

- I authorize the use of my protected health information as described above for the purpose(s) listed.
- I have the right to withdraw permission for the release of my information. If I sign this authorization to use or disclose information, I can revoke this authorization at any time. The revocation must be made in writing to the person/organization disclosing the information and will not affect information that has already been used or disclosed.
- I have a right to receive a copy of the authorization.

Unless revoked or otherwise indicated, the authorization's automatic expiration date will be one year from the date of my signature or upon the occurrence of the following event: _____

Signature of Patient/Legal Representative

Date

Description of Legal Representatives Authority

Appendix C

The Tennessee Department of Health School Accommodations Template for Concussion

Patient/Student: _____ Date: _____

Please excuse the above named patient from school today due to a medical appointment.

The student has sustained a concussion and is currently under the care of his or her physician and/or _____

the undersigned. S/he is not permitted to participate in any contact sport activity until formally cleared by his or her physician and/or the undersigned.

Please consider the following concussion-related recommendations:

Gym Class recommendations:

____ No gym class

____ Restricted gym class as specified: _____

Recommended **Academic** accommodations:

____ Untimed tests

____ Open note/open book or oral tests

____ Tutoring

____ Reduced workload when possible

____ 15 minute rest breaks from class every hour(s)

____ Modified/reduced homework assignments

____ Extended time on homework/projects

____ Tape record class lectures

____ Should not return to school until concussion symptoms are resolved

____ Other recommendations: _____

The patient/student will be re-evaluated on: _____

Healthcare Provider Name: _____ Address: _____

Signature: _____

Appendix D

CONCUSSION RETURN TO PLAY

Athlete's Name: _____ Date of Birth: _____

Date of Injury: _____

This return to play is based on today's evaluation Date of Evaluation: _____

Care Plan completed by: _____

Return to this office date/time: _____

Return to School date: _____

RETURN TO SPORTS INFO:

- 1 Athletes should not return to practice or play the same day that their injury occurred.
- 2 Athletes should never return to play or practice if they still have ANY symptoms – serious injury or death (although rare) can result
- 3 Athletes, be sure that your coach and/or athletic trainer are aware of your injury, symptoms and have the contact information for the health care provider treating your concussion.

Please initial:

_____ The athlete reports that he/she has no symptoms while participating in daily activities at this time.

_____ I have education the athlete and parents/guardian about the dangers of returning to play before symptoms have cleared.

The following are the return to sports recommendations at this time: (Please initial any recommendations selected)

PHYSICAL EDUCATION CLASS:

_____ Do NOT return to PE class at this time. (See "Return to this office date/time" above).

_____ Student MAY return to PE class after completion of Gradual Return to Play Plan (on back).

SPORTS:

_____ Do NOT return to sports practice or competition at this time.

_____ May GRADUALLY return to sports **activities** following the Gradual Return to Play Plan described on the back, under the supervision of the health care professional for your school or team.

_____ May be advanced back to **competition** after successful completion of the Gradual Return to Play Plan described on the back and after a **phone conversation** with treating health care provider.

_____ Must **return to the treating healthcare provider** for final clearance to return to competition after completing the Gradual Return to Play Plan. (See "Return to this office date/time" above).

_____ All steps of Return to Play Plan have been completed successfully. Cleared for full participation in all activities without restriction.

_____ No concussion suspected, cleared for full participation without a gradual return to play plan.

Appendix D

Treating Health Care Provider Information (Please print or stamp):

Provider's Name: _____ Provider's Office Phone: _____

Provider's Signature: _____ Office Address: _____

Please check:

Medical Doctor (MD) w/ concussion training

Osteopathic Physician (DO)

Clinical Neuropsychologist w/ concussion training

Physician Assistant (PA who is a member of a health care team supervised by a Tennessee licensed medical doctor or osteopathic physician.*

*Clearance by a PA is not accepted by the Tennessee Secondary School Athletic Association.

GRADUAL RETURN TO PLAY PLAN

Return to play should occur in gradual steps beginning with light aerobic exercise only to increase your heart rate (e.g. stationary cycle); moving to increasing your heart rate with movement (e.g. running); then adding controlled contact if appropriate; and finally return to sports competition.

Pay careful attention to your symptoms and your thinking and concentration skills at each stage of activity. After completion of each step **without recurrence of symptoms and no pain medication**, you can move to the next level of activity the next day. Move to the next level of activity only if you do not experience any symptoms at the present level. If your symptoms return, let your health care provider know, return to the first level of activity and restart the program gradually. This Gradual Return to Play process is for your own safety. Returning to play while still experiencing symptoms can result in serious injury or death. It is critical that you honestly report your symptoms to your doctor, coach and health care professional at the school.

GRADUAL RETURN TO PLAY PLAN:

"Day 1" means first day cleared to participate in Gradual Return to Play Plan, not first day after injury.

Day 1: Low levels of physical activity (i.e. symptoms do not come back during or after the activity). This includes walking, light jogging, light stationary biking and light weightlifting (low weight – moderate reps, no bench, no squats).

Day 2: Moderate levels of physical activity with body/head movement. This includes moderate jogging, brief running, moderate intensity on the stationary cycle, moderate intensity weightlifting (reduced time and or reduced weight from your typical routine).

Day 3: Heavy non-contact physical activity. This includes sprinting/running, high intensity stationary cycling, completing the regular lifting routine, non-contact sport-specific drills (agility with 3 planes of movement).

Day 4: Sports-specific practice.

Day 5: Full contact in a controlled drill or practice.

Day 6: Return to competition.

Adapted from the *Acute Concussion Evaluation Care Plan* from the Center for Disease Control and Prevention (<https://www.cdc.gov/injury/>), the *TSSAA Concussion Return to Play* form (<https://cms-files.tssaa.org/documents/tssaa/forms/Concussion-Return-to-Play-Form-updated-12.2019.pdf>) and the *TN Return to Learn/Return to Play: Concussion Management Guidelines*. All medical providers are encouraged to review the sites if they have questions regarding the latest information on the evaluation and care of a youth athlete following a concussion injury.

CDC'S ONLINE TRAINING FOR HEALTHCARE PROVIDERS

HEADS UP

HEADS UP to Healthcare Providers is a free online training developed by CDC and the American Academy of Pediatrics. The goal of the training is to provide an overview of the evidence-based recommendations outlined in the [CDC Pediatric mTBI Guideline](#) and to equip healthcare providers with practical strategies to integrate these recommendations into clinical practice.

WHAT YOU WILL LEARN

By the end of the training, you will be prepared to:


- ✦ Discuss what happens to the brain during and after an mTBI
- ✦ Identify at least three best practices related to diagnosis of mTBI
- ✦ Devise an appropriate management plan for pediatric patients with mTBI
- ✦ Describe prevention strategies for pediatric mTBI

FOLLOW THE URL TO BEGIN

[HTTPS://WWW.CDC.GOV/HEADSUP/PROVIDERS/TRAINING/](https://www.cdc.gov/headsup/providers/training/)



<https://www.tndisability.org/brain>



Research Summary and References

Support for the Toolkit

TOOLKIT

This toolkit, and specifically the *Concussion Management Protocol*, were developed based on the research summarized below. The research supports educating practitioners (rationale for the **Reference** section), properly evaluating, monitoring and referring patients (rationale for the **In-Office** section) and properly educating those with mTBI/ TBI (rational for the **Send-Home** sections).

CHILDREN:

Healthcare providers outside hospitals are on the front lines:

Most (82%) of those 0 to 17 years will seek initial care with their primary care physician (Arbogast, et al., 2016). Since most of our incidence data comes from Emergency Department's (ED's), we are significantly underestimating the extent of the TBI issue (Study included over 8,000 patients).

The very young are frequently not diagnosed or treated:

The newest pediatric mTBI guidelines recommend using an age-appropriate validated concussion scale (Lumba-Brown, et al., 2018), but one does not exist yet that focuses on children five and under. We must look for additional signs in children five years and under. For this age range, parents endorse the typical symptoms from the ACE, but in answer to an open-ended question, 82% also reported additional symptoms (Suskauer, et al., 2018), including:

- * Appetite changes
- * Behavioral dysregulation
- * Decreased engagement
- * Disrupted sleep
- * Bladder incontinence (Enuresis)
- * Increased dependence
- * Stomachaches

The study also concluded that it is important to monitor behavior dysregulation over time. At first, parents saw disengagement, and then behavior dysregulation emerged and persisted. Behavioral dysregulation was among most commonly reported symptoms and was still present at the time of the evaluation (over one month post).

Children with TBI may develop or have ongoing concerns and should be monitored (for years):

They are more likely to have a variety of health/academic issues compared to those with no TBI (Haarbauer-Krupa, Lee, et al., 2018). The highest prevalence are:

- * Learning disorders
- * ADD/ADHD
- * Speech Language problems
- * Developmental delay
- * Anxiety
- * Bone, joint or muscle problems

Children with mild (Taylor, 2015) and moderate and severe (Schwartz, 2003) injuries are more at risk for persistent behavior problems. The risk rises with severity of the mTBI and younger age at injury. Even in children whose injuries were significant enough to show skull or brain tissue damage on imaging, only one-fourth received any rehabilitations services afterward and only one-fourth received a neuropsychological assessment. None of the children received early intervention or special education preschool services after their TBI (Haarbauer-Krupa, Lundine, et al., 2018). This study concludes:

- * Healthcare providers should provide information to parents on what to watch for and long term implications.

- * Healthcare providers should make appropriate referrals at the time of diagnosis.
- * Referral to rehabilitation can help with transition to preschool.

Another study (Niedzwecki, et al., 2018) concluded that even though children did not receive inpatient care, some will still benefit from rehabilitation for subsequent problems, including memory and learning issues (that were not pre-existing).

This study also found that medical issues at the time of injury, like elevations or depressions of Intra-cranial pressure (ICP), unstable blood pressure, unstable oxygenation, delayed nutrition or seizures, can impact the child's IQ at 12 months.

- * The study's recommendation for trauma treatment is that rehab services be included early in the continuum – this would include consultation early in the ICU or acute care settings and referrals to an outpatient concussion clinic.

In the first year after injury, a substantial portion of children with moderate or severe TBI have unmet or unrecognized healthcare needs, with cognitive services being most frequent among these. Because of this finding, the authors recommended that cognition be screened in the primary care setting (Slomine, et al., 2006).

Reason for unmet needs:

- * Lack of a physician's recommendation or referral
- * Failure of parent follow-up
- * Not provided in the school settings
- * Cost

Children with all levels of impairment had educational needs, while those with less severe injuries were at greater risk of being underserved (Kingery, et al., 2017).

Earlier age at time of injury produces more functional impairment (Taylor, et al., 2015). The more severe the injury and the younger age at injury, the greater the need for monitoring and follow up (Anderson, Catroppa, Dudgeon, 2006; Anderson, Catroppa, Haritou, 2006).

On the first visit, provide educational materials, accommodations for return to school and recommend a follow up visit (at which time appropriate referrals can be made):

- * Many children did not even visit a healthcare provider in the year following their injury (Slomine, et al., 2006).

Ongoing family support is important:

Family support is important because those with family dysfunction/poor coping, the child had greater dysfunction (Schwartz, 2003; Anderson, Catroppa, Dudgeon, et al., 2006; Taylor, 2008).

Families also reported needing information, emotional support and access to community-based services (Jones, 2017).

Schools need the support/recommendations of healthcare providers:

Teachers are not adequately trained to identify brain injuries and issues related to them (Davies, et al., 2013).

On specialized testing, children with TBI tend to show specific patterns of deficit that will not be revealed through standard special education testing. A neuropsychological evaluation will pick up these patterns. In a study of mild complicated TBI (with orthopedic controls), children who were injured before age 6 and were about 5 years post injury were tested. Both groups were within normal limits on most cognitive, language and reading measures; but they had some differences in verbal IQ, receptive

language and reading comprehension. The biggest differences were in pragmatic language (which leads to social issues), story retell, and word fluency (Haarbauer-Krupa, King, et al., 2019).

Schools will not provide all of what a child needs (Niedzwecki, 2018). Schools are only required to provide those services that directly relate to academics.

The gap in academic achievement widens over time (compared with non-injured classmates) (Ewing-Cobbs, 2006; Farmer, 1997; Taylor & Yeates, 2002; Todis & Glang, 2008; Todis, Glang, Bullis, et al., 2011; Wagner, et al., 2006). So, if children with TBI do not qualify for services at first, they should be referred again if they continue to have difficulties.

“Children who receive systematic transition services a part of their medical care are more likely to be identified for specialized support services at school, such as speech therapy (Haarbauer-Krupa, Ciccio, et al., 2017).

Use of the ACE tools (screening tool and Care Plan) “increased patient follow-up and improved recall of and adherence to ED discharge recommendations (Zuckerbraun, 2014).”

Pediatric Guideline:

Also see the CDC Pediatric Guideline (Lumba-Brown, et al., 2018) on mTBI in this toolkit for 19 sets of recommendations, with these [5 key take away points](#):

1. Do not routinely image pediatric patients to diagnose mTBI.
2. Use validated, age-appropriate symptom scales to diagnose mTBI.
3. Assess risk factors for prolonged recovery, including history of mTBI or other brain injury, severe symptom presentation immediately after the injury, and personal characteristics and family history (such as learning difficulties and family and social stressors).
4. Provide patients and their parents with instructions on returning to activity customized to their symptoms.
5. Counsel patients and their parents/caregivers to return gradually to non-sports activities after no more than 2-3 days of rest.

Consequences of brain injury for all ages:

Once a person has one brain injury, the risk for another increases, and the risk increases with each subsequent injury. A person with a brain injury is also more likely to be incarcerated (or involved with the criminal justice system) (Farrer & Hedges, 2011; Shiroma, et al., 2012; Williams, et al., 2010; Im, et al., 2014), to have psychiatric issues ((McCarthy, et al., 2006; Kaponen, et al., 2002; Zgaljardic, et al., 2015), to be involved with substance abuse (Kreutzer, et al., 1996), and to be socially isolated (Morton & Wehman, 1995; Hawthorne, et al., 2009). Long-term psychiatric disorders are associated with greater risk for substance abuse (Zgaljardic, et al., 2015). Prior TBI has been identified as a potential contributing factor to domestic violence (Romero-Martinez & Moya-Albiol, 2013). Not surprisingly, TBI is found in female victims of domestic violence (Corrigan, et al., 2001).

ADULTS

Follow up and education are important:

Findings from a study (Seabury, et al., 2018) of follow-up care that was provided to people at 11 Level 1 trauma centers across the country:

- * Less than half received TBI educational material at discharge or saw a health care practitioner within 3 months after injury.
- * Only 27% were called by 2 weeks.
- * Follow-up care varied by site, from 19% to 72%.

- * For those with a positive CT scan, over one-third had not seen a medical practitioner for follow-up.
- * Even among those with 3 or more moderate to severe post-concussive symptoms, only about half saw a medical practitioner within 3 months.
 - o Of those that did, 80% reported that it was helpful. The majority saw a general practitioner and 38% saw a neurologist. Only 15% reported visiting a clinic specializing in TBI care.

A few conclusions from the paper:

- * “Failure to follow-up with patients could have adverse consequences, as simply providing educational materials to patients with mTBI is associated with improved outcomes.”³⁵
- * “Our findings reveal the consequences that may result from the absence of systems of follow-up care for patients with mTBI and concussion. They also highlight an apparent lack of appreciation by many clinicians of the substantial symptom and life burdens experienced by a significant proportion of patients with injuries labeled mild.”

Use of the ACE tools (screening tool and Care Plan) “increased patient follow-up and improved recall of and adherence to ED discharge recommendations (5-21 year olds) (Zuckerbraun, 2014).”

Unmet Needs:

Poor psychosocial health was reported by a substantial portion in a study at one year post injury TBI may cause decades lasting vulnerability to psychiatric illness in some individuals. They were most susceptible to depression, delusional disorders and personality disturbances. This study highlights the importance of psychiatric follow up even decades (30 years) later (Kaponen, et al., 2002). Heinemann found unmet needs at 7 years. The most prevalent were improving memory and problem solving, increasing income and improving job skills (Heinemann, et al., 2002).

[Also see the Updated Mild Traumatic Brain Injury Guideline for Adults in this toolkit.](#)

Model of 6 types of concussion and active treatments (pediatric and adult):

There is now a great body of evidence supporting the 6 types of concussion and the active treatments for each type. A good resource to start with is *Concussion: A Clinical Profile Approach to Assessment and Treatment* by Kontos and Collins (2018) and *A comprehensive, targeted approach to the clinical care of athletes following sport-related concussion* (Collins, et al., 2013).

References

- Anderson, V. A., Catroppa, C., Dudgeon, P., Morse, S. A., Haritou, F., & Rosenfeld, J. V. (2006). Understanding predictors of functional recovery and outcome 30 months following early childhood head injury. *Neuropsychology*, 20(1), 42-57. doi:10.1037/0894-4105.20.1.42
- Anderson, V., Godfrey, C., Rosenfeld, J.V., & Catroppa, C., Predictors of Cognitive Function and Recovery 10 Years After Traumatic Brain Injury in Young Children. (2012). *Pediatrics*, 129(2). doi:10.1542/peds.2011-0311d
- Arbogast, K. B., Curry, A. E., Pfeiffer, M. R., Zonfrillo, M. R., Haarbauer-Krupa, J., Breiding, M. J., . . . Master, C. L. (2016). Point of Health Care Entry for Youth With Concussion Within a Large Pediatric Care Network. *JAMA Pediatrics*, 170(7). doi:10.1001/jamapediatrics.2016.0294
- Collins, M. W., Kontos, A. P., Reynolds, E., Murawski, C. D., & Fu, F. H. (2013). A comprehensive, targeted approach to the clinical care of athletes following sport-related concussion. *Knee Surgery, Sports Traumatology, Arthroscopy*, 22(2), 235-246. doi:10.1007/s00167-013-2791-6
- Corrigan, J. D., Wolfe, M., Mysiw, W. J., Jackson, R. D., & Bogner, J. A. (2001). Early identification of mild traumatic brain injury in female victims of domestic violence. *Clinical Journal of Womens Health*, 01(4), 184-190. doi:10.1053/cjwh.2001.27867

- Davies, S. C., Fox, E. E., Glang, A., et al. Traumatic Brain Injury and Teacher Training: A Gap in Educator Preparation. (2013). Counselor Education and Human Services Faculty Publications. 25. http://ecommons.udayton.edu/edc_fac_pub/25
- Ewing-Cobbs, L., Prasad, M. R., Kramer, L., Cox, C. S., Baumgartner, J., Fletcher, S., et al. (2006). Late intellectual and academic outcomes following traumatic brain injury sustained during early childhood. *Journal of Neurosurgery: Pediatrics*, 105(4), 287-296. doi:10.3171/ped.2006.105.4.287
- Farmer, J. E., & Johnson-Gerard, M. (1997). Misconceptions about traumatic brain injury among educators and rehabilitation staff: A comparative study. *Rehabilitation Psychology*, 42(4), 273-286. doi:10.1037//0090-5550.42.4.273
- Farrer, T. J., & Hedges, D. W. (2011). Prevalence of traumatic brain injury in incarcerated groups compared to the general population: A meta-analysis. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 35(2), 390-394. doi:10.1016/j.pnpbp.2011.01.007
- Haarbauer-Krupa, J., King, T. Z., Wise, J., Gillam, S., Trapani, J., Weissman, B., & Depompei, R. (2019). Early Elementary School Outcome in Children With a History of Traumatic Brain Injury Before Age 6 Years. *Journal of Head Trauma Rehabilitation*, 34(2), 111-121. doi:10.1097/htr.000000000000041
- Haarbauer-Krupa, J., Lee, A. H., Bitsko, R. H., Zhang, X., & Kresnow-Sedacca, M. (2018). Prevalence of Parent-Reported Traumatic Brain Injury in Children and Associated Health Conditions. *JAMA Pediatrics*, 172(11), 1078. doi:10.1001/jamapediatrics.2018.2740
- Haarbauer-Krupa, J., Lundine, J. P., Depompei, R., & King, T. Z. (2018). Rehabilitation and school services following traumatic brain injury in young children. *NeuroRehabilitation*, 42(3), 259-267. doi:10.3233/nre-172410
- Hawthorne G, Gruen RL, Kaye AH. Traumatic brain injury and long-term quality of life: findings from an Australian study. *J Neurotrauma*. 2009;26:1623–33. doi: 10.1089/neu.2008.0735
- Heinemann, A. W., Sokol, K., Garvin, L., & Bode, R. K. (2002). Measuring unmet needs and services among persons with traumatic brain injury. *Archives of Physical Medicine and Rehabilitation*, 83(8), 1052-1059. doi:10.1053/apmr.2002.3428
- Im, B., Hada, E., Smith, M., & Gertisch, H. (2014). The Relationship between TBI and incarceration rates. *Spotlight on Disability Newsletter*, Dec 2014 retrieved from <https://www.apa.org/pi/disability/resources/publications/newsletter/2014/12/incarceration>.
- Jones, S., Davis, N., & Tyson, S. F. (2017). A scoping review of the needs of children and other family members after a child's traumatic injury. *Clinical Rehabilitation*, 32(4), 501-511. doi:10.1177/0269215517736672
- Kaponen, S., Taiminen, T., Portin, R., et al. (2002). Axis I and II psychiatric disorders after traumatic brain injury a 30 year follow-up study. *Am J Psychiatry*, Aug; 159(8): 1315-21.
- Kontos, A. P., & Collins, M. W. (2018). *Concussion: A clinical profile approach to assessment and treatment*. Washington, DC: American Psychological Association.
- Kreutzer, J. S., Witol, A. D., & Marwitz, J. H. (1996). Alcohol and Drug Use Among Young Persons with Traumatic Brain Injury. *Journal of Learning Disabilities*, 29(6), 643-651. doi:10.1177/002221949602900608
- Lumba-Brown, A., Yeates, K.O., Sarmiento, K., et al. Centers for Disease Control and Prevention guideline on the diagnosis and management of mild traumatic brain injury among children [published online September 4, 2018]. *JAMA Pediatr*. doi:10.1001/jamapediatrics.2018.2853
- Mccarthy, M. L., Dikmen, S. S., Langlois, J. A., Selassie, A. W., Gu, J. K., & Horner, M. D. (2006). Self-Reported Psychosocial Health Among Adults With Traumatic Brain Injury. *Archives of Physical Medicine and Rehabilitation*, 87(7), 953-961. doi:10.1016/j.apmr.2006.03.007
- Morton, M. V., & Wehman, P. (1995). Psychosocial and emotional sequelae of individuals with traumatic brain injury: A literature review and recommendations. *Brain Injury*, 9(1), 81-92. doi:10.3109/02699059509004574
- Niedzwecki, C. M., Rogers, A. T., & Fallat, M. E. (2018). Using Rehabilitation along the Pediatric Trauma Continuum as a Strategy to Define Outcomes in Traumatic Brain Injury. *Clinical Pediatric Emergency Medicine*, 19(3), 260-271. doi:10.1016/j.cpem.2018.08.005

- Romero-Martinez, A., Moya-Albiol, L., Neuropsychology of perpetrators of domestic violence: the role of traumatic brain injury and alcohol abuse and/or dependence. (2013). *Revista de Neurologica*, Dec; 57(11): 515-522.
- Schwartz, L. (2003). Long-Term Behavior Problems Following Pediatric Traumatic Brain Injury: Prevalence, Predictors, and Correlates. *Journal of Pediatric Psychology*, 28(4), 251-263. doi:10.1093/jpepsy/jsg013
- Seabury, S.A., Gaudette, E., Goldman, A. J., et al. (2018). Assessment of Follow-up Care After Emergency Department Presentation for Mild Traumatic Brain Injury and Concussion: Result from the TRACK-TBI Study. *JAMA Network Open*. 208;1(1) doi:10.1001/jamanetworkopen.2018.0210
- Shiroma, E. J., Ferguson, P. L., & Pickelsimer, E. E. (2012). Prevalence of Traumatic Brain Injury in an Offender Population. *Journal of Head Trauma Rehabilitation*, 27(3). doi:10.1097/htr.0b013e3182571c14
- Slomine, B. S., McCarthy, M.L., Ding, R., et al. (2006). Health Care Utilization and Needs After Pediatric Traumatic Brain Injury. *Pediatrics*, 117(4). doi:10.1542/peds.2005-1892
- Suskauer, S. J., Rane, S., Reesman, J., & Slomine, B. S. (2018). Caregiver-report of symptoms following traumatic brain injury in a small clinical sample of preschool-aged children. *Journal of Pediatric Rehabilitation Medicine*, 11(1), 7-14. doi:10.3233/prm-160424
- Taylor, H. G., Swartwout, M. D., Yeates, K. O., Walz, N. C., Stancin, T., & Wade, S. L. (2008). Traumatic brain injury in young children: Postacute effects on cognitive and school readiness skills. *Journal of the International Neuropsychological Society*, 14(5), 734-745. doi:10.1017/s1355617708081150
- Taylor, H.G., Orchinik, L.J., Minich, N., et al. (2015). Symptoms of Persistent Behavior Problems in Children with Mild Traumatic Brain Injury. *J Head Trauma Rehabil*. Sep-Oct;30(5):302-10. doi:10.1097/HTR.000000000000106
- Taylor, H.G., Yeates, K. O., Wade, S.L., et al. (2002). A prospective study of short- and long-term outcomes after traumatic brain injury in children: Behavior and achievement. *Neuropsychology*, 16(1) 15-27.
- Todis, B., & Glang, A. (2008). Redefining Success. *Journal of Head Trauma Rehabilitation*, 23(4), 252-263. doi:10.1097/01.htr.0000327257.84622.bc
- Todis, B., Glang, A., Bullis, M., Ettel, D., & Hood, D. (2011). Longitudinal Investigation of the Post-High School Transition Experiences of Adolescents With Traumatic Brain Injury. *Journal of Head Trauma Rehabilitation*, 26(2), 138-149. doi:10.1097/htr.0b013e3181e5a87a
- Updated Mild Traumatic Brain Injury Guideline for Adults, retrieved from https://www.cdc.gov/traumaticbraininjury/mtbi_guideline.html
- Wagner, M., Newman, L., Cameto, R., Levine, P., and Garza, N. (2006). An Overview of Findings From Wave 2 of the National Longitudinal Transition Study-2 (NLT2). (NCSE 2006-3004). Menlo Park, CA: SRI International.
- Williams, W. H., Mewse, A. J., Tonks, J., Mills, S., Burgess, C. N., & Cordan, G. (2010). Traumatic brain injury in a prison population: Prevalence and risk for re-offending. *Brain Injury*, 24(10), 1184-1188. doi:10.3109/02699052.2010.495697
- Zgaljardic DJ, Seale GS, Schaefer LA, Temple RO, Foreman J, Elliott TR. Psychiatric disease and post-acute traumatic brain injury. *J Neurotrauma*. 2015;32:1911–25. doi: 10.1089/neu.2014.3569
- Zuckerbraun NS, Atabaki,S, Collins, MW, Thomas D, Gioia GA. Use of modified acute concussion evaluation tools in the emergency department. *Pediatrics*, 133(4). doi: 10.1542/peds.2013-2600d



www.tndisability.org/brain




Brain Links is supported by the Administration for Community Living (ACL) of the U.S. Department of Health and Human Services under Grant No. 90TBSG0024-01-00 and in part by the TN Department of Health, Traumatic Brain Injury Program.



TRAUMATIC BRAIN INJURY/ CONCUSSION

THINGS TO WATCH FOR OVER TIME:

- Headaches
- Changes in sleep patterns
- Fatigue
- Changes in vision
- Balance, coordination changes, dizziness
- Mood swings, gets mad easily
- Changes in personality
- Not feeling like themselves
- Trouble with attention and thinking
- Memory problems, especially short term
- Depression/Anxiety
- Difficulty handling stress
- Innapropriate behavior
- Grades dropping, falling behind in class
- Changes in work performance



Thank You!

We're here to help

Our mission is to bring together professionals to recognize the far-reaching and unique nature of brain injury and to improve services for survivors. If we can help you, please feel free to reach out!



Contact us:

tbi@tndisability.org

Check out our website:

www.tndisability.org/brain

Follow us on social media:



We want to hear from you!



Complete our short survey to let us know how we're doing.

