Brain Injury and Chronic Pain



Pain is the most common chronic medical condition reported by people with Traumatic Brain Injury (TBI).

Over 50%



Over 50% of people with TBI develop chronic pain



11 Times



People with TBI are at eleven times greater risk of accidental overdose

Common problems following brain injury, like poor judgment, memory and increased impulsivity make it harder to self-regulate substance use and make overdose more likely.

BEST PRACTICE Providers
specializing in
treating Chronic
Pain should:

- **SCREEN** for prior history of Brain Injury
- ASSESS Cognitive & Functional Impairment
- EDUCATE staff on Brain Injury
- EDUCATE the person about their Brain Injury
- PROVIDE and TEACH Accommodations
- **CONNECT** person served with Community Resources

COMMON PROBLEMS

After Brain Injury, we often see problems with:

- Attention, memory and new learning
- Slowed speed of processing
- Organization, problem solving & impulsivity
- Irritability, frustration & agitation
- Balance, dizziness & headaches
- Poor awareness of deficits & difficulties
- Difficulty being flexible, poor self-monitoring

WHAT TO LOOK FOR Providers specializing in treating Chronic Pain may see:

- Looking uninterested because they cannot pay attention
- Missing appointments
- Appearance of non-compliance because they cannot remember dosages and medication schedules
- Slow to follow directions because they cannot process quickly
- Falling into things, often getting hurt
- Difficulty re-entering community because of cognitive changes
- Gets stuck on an idea or a way of doing something, does not recognize mistakes

An estimated 50 million adults in the United States experienced chronic pain (i.e., pain lasting ≥3 months) in 2016, resulting in substantial health care costs and lost productivity. - CDC.gov

Common Accommodations for Brain Injury Challenges

Here are some common and simple accommodations:

- Working for shorter periods of time
- o Getting rid of distractions around you, like noise or movement
- Taking notes (on paper, in a notebook, on a phone or computer)
- Using a phone to set timers to remember appointments & medication schedules; providing a check-off medication schedule to avoid forgetting they have already taken a medication
- Repeating information to the person
- Slowing down when talking; giving them more time to respond
- o Giving a written list of non-medication strategies to avoid or reduce pain
- Coaching the person to take deep breaths when feeling angry or anxious

Tools for Best Practice

Brain Links' Strategies & Accommodations Tool: https://www.tndisability.org/rehabilitation
Symptom Questionnaire: https://mindsourcecolorado.org/adult-symptom-questionnaire/
Cognitive Strategies Guidebook: https://mindsourcecolorado.org/wpcontent/uploads/2019/05/Strategies-and-Accommodations-Guidebook-CJ-Professionals-5.6.19.pdf
Brain Injury Screening Resources:

- OSU TBI Identification Method: https://wexnermedical.osu.edu/neurological-institute/neuroscience-research-institute/research-centers/ohio-valley-center-for-brain-injury-prevention-and-rehabilitation/osu-tbi-id
- NASHIA's OBISSS: https://www.nashia.org/obisssprogram

Tennessee Resources

<u>Brain Links' Website</u> with many resources: tndisability.org/brain

<u>Brain Links' Toolkits</u> (for Service Professionals and Survivors): https://www.tndisability.org/brain-toolkits

TN Department of Health TBI Program:

https://tinyurl.com/3v5jrdt3



<u>Training for Brain Injury:</u>
https://www.tndisability.org/tbf-brain-injury



Resource Pages & Training for Chronic Pain: https://www.tndisability.org/tbf-pain

TN Brighter Futures is organized and facilitated by Brain Links through a contract from the TN Department of Health TBI Program.





Pain Resources

References:

- Hammond, F. M., Barrett, R. S., Shea, T., et. al. (2015). Psychotropic medication use during inpatient rehabilitation for traumatic brain injury. Archives of Physical Medicine and Rehabilitation, 96 (8Suppl 3): S256-73. 16.
- Hammond, F. M., Dams-O'connor, K, Ketchum J et al (2018). Mortality secondary to accidental poisoning after inpatient rehabilitation for traumatic brain injury study: A NIDILRR Traumatic Brain Injury Model Systems Study.
- John D. Corrigan and Rachel Sayko Adams (2019). The Intersection of Lifetime History of Traumatic Brain Injury and the Opioid Crisis. Addictive Behaviors; 90: 143–145. https://doi.org/10.1016/j.addbeh.2018.10.030









